

---

## Use of botulinum toxin in surgical preparation for complex incisional hernias

---

João Kleber de Almeida Gentile 1, Amanda Salvati Beck Lima 2, Cassiana de Moura e Costa 2, Gabriella Muraro 2, Lorena de Freitas da Silva 2

<sup>1</sup> Médico Cirurgião do Aparelho Digestivo. Membro Titular Especialista do Colégio Brasileiro de Cirurgia Digestiva (CBCD). Fellow of American College of Surgeons (FACS) Professor da Disciplina de Técnica Operatória e Habilidades Cirúrgicas da Faculdade de Medicina da Universidade Cidade de São Paulo (FM-UNICID). São Paulo, SP, Brazil.

<sup>2</sup> Aluno(a) de Ciências Médicas da Faculdade de Medicina do Centro Universitário Assis Gurgacz. Cascavel, PR, Brazil.

\*Corresponding author: João Kleber de Almeida Gentile. Rua Padre Estevão Pernet, 180 – Tatuapé. Zip Code: 03315-000 – São Paulo, SP, Brazil. Phone: +55 (11) 2093-1303. E-mail: joaokleberg@gmail.com.

**Research Ethics Committee Approval (if necessary):** We declare that the patient approved the study by signing an informed consent form and the study followed the ethical guidelines established by the Declaration of Helsinki.

Received on: Apr 14, 2022. Accepted on: Jun 11, 2022. Available online: Jun 13, 2022.

---

### Abstract

Abdominal wall defects with incisional hernias after surgery represent a great challenge for surgeons due to their technical complexity for treatment, largely attributed to the retraction of the lateral muscles of the abdominal wall. Reconstruction of the abdominal wall is understood to re-functionalize it with the approximation of the rectus abdominis muscles through its midline. The use of botulinum toxin serotype A (TBA) has shown that its use produces a temporary flaccid paralysis in the abdominal muscles, favoring midline closure, caused by muscle flaccidity and greater complacency caused by the toxin. Abdominal hernias result in increased morbidity due to lung restriction, aerophagia, abdominal pain, constipation, and urinary alterations), with a consequent decrease in quality of life. The objective of this study is describe our initial experience with chemical separation of components using the TBA in Brazil.

**Keywords:** Incisional Hernia; Botulinum Toxins, Type A; Abdominal wall.

### Introduction

Incisional hernia is a common type of hernia of the abdominal wall at

the site of an anterior surgical incision, with a prevalence of more than 30% in high-risk groups and is most often

located in the midline of the abdomen, resulting from a failure of abdominal wall closure [1-3]. This complication has been shown to generate a considerable cost to society, as hernia-related expenses in the United States increased by 52% from 2007 to 2011, estimated to be approximately \$7.3 billion annually [4-6]. As a result of the challenges of abdominal wall surgeries, studies have demonstrated the use of botulinum toxin A (TBA) injection to aid in the operative preparation of incisional hernias [7, 8].

TBA is a neurotoxin that acts selectively on the presynaptic cholinergic nerve terminals, blocking the release of acetylcholine. This action will promote relaxation of the lateral muscles of the abdominal wall (internal oblique, external oblique, and transverse abdominal), facilitating the closure of the aponeurosis, as well as decreasing postoperative pain [8-10].

The goal of using TBA is to perform an elective incisional hernia preparation, to prevent early recurrence and decrease the risk of abdominal compartment syndrome and some complications such as incarceration, bowel obstruction, and incarceration. In this manuscript, we relate the initial experience of our institution with the use of TBA in the surgical preparation for wall reconstructions in patients with complex abdominal wall hernias.

## Experience Report

The abdominal wall plasty is understood to refunctionalize the anatomy and function of the abdominal wall with the approximation of the rectus abdominis muscles through its midline. The use of TBA has demonstrated that its use produces a temporary flaccid paralysis in the abdominal muscles, favoring the closure of the midline, caused by muscular flaccidity and greater complacency provoked by the toxin [2, 4].

The objective of the application of TBA in the preoperative in patients with incisional hernias with defects larger than 10 cm, as an alternative to the treatment with pneumoperitoneum, in face of its high rate of complications such as infection, hospital stay, sometimes longer than 15 days, and complexity (higher cost than the use of botulinum toxin).

### Application Protocol:

1. Patients with giant incisional hernia with loss of domicile (>10 cm) and indication for surgical treatment without clinical contraindication;
2. Evaluation with computed tomography (CT) of abdomen and pelvis with measurement of abdominal volumetry;
3. Botulinum toxin A (TBA) injection;

- a. Interventional radiology (puncture guided ultrasonography);
- b. 5 puncture points according to Figure 1;
- c. 1 ampoule (100UI) diluted in 2 mL of sterile solution (total of 10UI in each point);
- d. Reevaluation by CT of abdomen and pelvis in 21 days for new abdominal volumetry.

In our surgical service we currently have 4 patients who underwent TBA guided by ultrasonography respecting the indications in the literature with satisfactory results from the surgical point of view, without any clinical or surgical complication related to the technique described.



**Figure 1.** Points of preoperative botulinum toxin infiltration, showing 2 points in the medial axillary line and 3 points in the external oblique muscle guided by ultrasound.

Despite being cases of complex hernias, the use of TBA showed to be efficient in the surgical preparation of these patients, leading the cases to an

easier aponeurotic and fascial closure, without presenting recurrence of the cases as described in table 1.

**Table 1.** Analysis and follow-up of patients submitted to the use of TBA in the surgical preparation of complex incisional hernia repair.

Years	BMI (Kg/m <sup>2</sup> )	Size of the Defect Diameter on the Largest Shaft (mm )	Complications	Follow-up (months)
57	29,2	110	No	4
44	26,4	125	No	8
61	24,9	140	No	11
59	31,1	120	No	12

## Discussion and Conclusion

Incisional hernias occur in 15-28% of all patients undergoing abdominal surgery, and their repair is a complex procedure with high failure rates. It is estimated that within three years the hernia recurrence rate ranges from 15-21% for open repair and 7-15% for laparoscopic repair. This vicious cycle of recurrence contributes to worse surgical outcomes for patients in addition to greater expense to the healthcare system [ 11-13].

TBA is a neurotoxin isolated and purified from *Clostridium botulinum* type A bacteria and acts at the neuromuscular junction by inhibiting the release of acetylcholine and causing flaccid paralysis [10, 14]. The effects of botulinum toxin begin to appear within two to three days, with complete effect in two weeks and decline after 3 months. Application of this toxin to the lateral musculature (internal oblique, external oblique, and transverse abdominis) can paralyze and subsequently lengthen the muscle complex.

Thus, facilitating the medialization of the rectus muscles and the closure of the abdominal wall [15, 16].

Furthermore, several research have proven that the use of TBA in the preoperative management of hernias did not bring significant complications during and after the surgical procedure. A clinical trial, published in 2020, compared 80 consecutive patients with large midline incisional hernias. The patients were divided into 2 groups, of which 40 had preoperative TBA administration and open Rives repair, and the rest underwent only open component separation during the observation period. The result of the prospective, comparative study was that complete fascial closure was possible in all subjects in the BTA group and that there were no complications during the administration of the neurotoxin in the preoperative period [17].

A systematic review published in 2017 demonstrated that in addition to the importance of BTA in hernia repair

favoring fascial closure, long-term analgesic effects were also obtained through ultrasound injection of Botox and could relieve pain for up to 3 months. Moreover, use of Botox preoperatively reduced the width of the ventral hernia and lengthening of the abdominal muscles bilaterally, thus favoring the approximation of the borders without interrupting healing [18].

Protocols for the use of TBA have been described in the literature, in which the injection of the toxin was always guided by an imaging exam (tomography, electromyography and ultrasonography), with ultrasonography being the most used. The injection should be applied at least 2 weeks before reconstruction of the abdominal wall, with maximum effect between 2 and 4 weeks.

The most used botulinum toxin dosages were 200-300 units of TBA-Botox® or 500 units of TBA-Dysport® in a 2:1 saline dilution, with Dysport® being less potent. In all studies, both the external and internal oblique muscles were injected, however the vast majority of studies BTX was injected into all 3 muscles (internal oblique, external oblique and transverse abdominis) [19].

Also, Ibarra-Hurtado et al. [11] published two studies, which included all 29 patients with midline defects. They received a total dose of 500 units of

TBA 4 weeks before the hernia repair. The surgical procedure was performed with either retro muscular mesh placement or through simple suture closure without mesh. Successful primary fascial closure was achieved in 100% of patients, however, 15 patients (53%-23 and 50%-24) still required additional component separation techniques for abdominal closure [19, 20].

The use of TBA, besides contributing to the flaccid paralysis of muscular tissues and relaxation of the abdominal wall, facilitating the repair of complex hernias, there still seems to be a gain of tissue, even if partial, increasing the chance of primary closure of the abdominal wall, without the need for bridging closure or complex and expensive meshes for closure. In our experience we could perceive an easier abdominal wall closure, without presenting any related complications such as abdominal compartment syndrome or local complications. It is evident that the use of TBA can be a great help in the treatment of complex ventral hernias.

Use of TBA has been widely used in the treatment of complex hernias of the abdominal wall, being increasingly indicated against other more expensive and less safe options. More studies and systemic reviews are still needed to evaluate the benefit of its long-term use

and consolidate the indication of TBA as a routine in the surgical preparation of these patients with large abdominal wall defects.

## References

- [1] Hope WW, Tuma F. Incisional Hernia. 2021 Jul 18. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2022 Jan–.
- [2] Berrevoet F. Prevention of Incisional Hernias after Open Abdomen Treatment. *Front Surg*. 2018 Feb 26;5:11. doi: 10.3389/fsurg.2018.00011.
- [3] Um P Jairam. Timmermans L. Eker Hasan H. Prevention of Incisional hernia with prophylactic onlay and sublay mesh reinforcement versus primary suture only in midline laparotomies (PRIMA): 2-year follow-up of a multicentre, double-blind, randomised controlled trial. *Lancet*. 2017. Aug 5. 390(10094):567-576. doi: 10.1016/S0140-6736(17)31332-6.
- [4] Bower C, Roth JS. Economics of abdominal wall reconstruction. *Surg Clin North Am*. 2013 Oct;93(5):1241-53. doi: 10.1016/j.suc.2013.06.007.
- [5] Reynolds D, Davenport DL, Korosec RL, Roth JS. Financial implications of ventral hernia repair: a hospital cost analysis. *J Gastrointest Surg*. 2013 Jan;17(1):159-66; discussion p.166-7. doi: 10.1007/s11605-012-1999-y.
- [6] Basta MN, Kozak GM, Broach RB, Messa CA 4th, Rhemtulla I, DeMatteo RP, Serletti JM, Fischer JP. Can We Predict Incisional Hernia?: Development of a Surgery-specific Decision-Support Interface. *Ann Surg*. 2019 Sep;270(3):544-553. doi: 10.1097/SLA.0000000000003472.
- [7] Walming S, Angenete E. Retrospective review of risk factors for surgical wound dehiscence and incisional hernia. *BMC Surg*. 2017. Feb 22. 17(1):19. doi: 10.1186/s12893-017-0207-0.
- [8] Rosin D. Outcomes of Chemical Component Paralysis Using Botulinum Toxin for Incisional Hernia Repairs. *World Journal of Surgery*. 2013. Oct 1; 37 (12):2838-8.
- [9] Whitehead Clarke T, Windsor A. The Use of Botulinum Toxin in Complex Hernia Surgery: Achieving a Sense of Closure. *Frontiers in Surgery*. 2021. Oct 1;8.
- [10] Mourad AP, De Robles MS, Winn RD. Low Dose Pre Operative Botulinum Toxin A Effectively Facilitates Complex Ventral Hernia Repair: A Case Report and Review of the Literature. *Medicina (Kaunas)*. 2020. Dec 28; 57 (1):14.
- [11] Ibarra-Hurtado TR, Nuño-Guzmán CM, Miranda-Díaz AG, Troyo-Sanromán R, Navarro-Ibarra R, Bravo-Cuéllar L. Effect of botulinum toxin type A in lateral abdominal wall muscles

thickness and length of patients with midline incisional hernia secondary to open abdomen management. *Hernia*. 2014 Oct;18(5):647-52.

[12] Bueno-Lledó J, Torregrosa A, Ballester N, Carreño O, Carbonell F, Pastor PG, Pamies J, Cortés V, Bonafé S, Iserte J. Preoperative progressive pneumoperitoneum and botulinum toxin type A in patients with large incisional hernia. *Hernia*. 2017 Apr;21(2):233-243. doi: 10.1007/s10029-017-1582-2.

[13] Bueno-Lledó J, Torregrosa Gallud A, Jiménez Rosellón R, Carbonell Tatay F, García Pastor P, Bonafé Diana S, Iserte Hernández J. Preoperative preparation of «loss of domain» hernia. Progressive pneumoperitoneum and botulinum toxin type A. *Cir Esp*. 2017 May;95(5):245-253. English, Spanish. doi: 10.1016/j.ciresp.2017.04.

[14] Nigam PK, Nigam A. Botulinum toxin. *Indian J Dermatol*. 2010;55(1):8-14. doi: 10.4103/0019-5154.60343.

[15] Farooque F, Jacombs AS, Roussos E, Read JW, Dardano AN, Edye M, Ibrahim N. Preoperative abdominal muscle elongation with botulinum toxin A for complex incisional ventral hernia repair. *ANZ J Surg*. 2016 Jan-Feb;86(1-2):79-83. doi: 10.1111/ans.13258.

[16] Whitehead-Clarke T, Windsor A. The Use of Botulinum Toxin in Complex

Hernia Surgery: Achieving a Sense of Closure. *Front Surg*. 2021; 8: 753889. doi: 10.3389/fsurg.2021.753889.

[17] Bueno-Lledó J, Martinez-Hoed J, Torregrosa-Gallud, Menéndez-Jiménez M, Pous-Serrano S. Botulinum toxin to avoid component separation in m midline large hernias. *Surg Journal*. 2020. Vol 168, Issue 3, P543-549.

[18] Weissler J M, Lanni M A, Tecce M G, Carney M J, Shubinets V, Fischer J P. Chemical component separation: a systematic review and meta-analysis of botulinum toxin for management of ventral hernia. *Journal of plastic surgery and hand surgery*. 2017. doi: 10.1080/2000656X.2017.1285783.

[19] Ibarra-Hurtado TR, Nuño-Guzmán CM, Miranda-Díaz AG, Troyo-Sanromán R, Navarro-Ibarra R, Bravo-Cuellar L. Effect of botulinum toxin type A in lateral abdominal wall muscles thickness and length of patients with midline incisional hernia secondary to open abdomen management. *Hernia*. 2014 Oct;18(5):647-52.

[20] Ibarra-Hurtado TR, Nuño-Guzmán CM, Echeagaray-Herrera JE, Robles-Vélez E, de Jesús González-Jaime J. Use of botulinum toxin type a before abdominal wall hernia reconstruction. *World J Surg*. 2009 Dec;33(12):2553-6. doi: 10.1007/s00268-009-0203-3.

**Conflict of interest:** The author declares no conflicts of interest.

**Acknowledgements:** None.

**Funding:** None.

**How to cite this article:** Gentile JKA, Lima ASB, Costa CM, Muraro G, da Silva LFS. Use of botulinum toxin in surgical preparation for complex incisional hernias. Brazilian Journal of Case Reports. 2022 Jul-Sep;02(3):75-82.